


REMARKS

By the present Amendment, claims 1-9 are amended in accordance with customary U.S. patent practice and to omit the multiple dependency of the claims. A Version With Markings Showing Changes Made is attached. Claims 10-20 are added. Support for claims 10, 11, and 12 and 13 may be found in original claims 3, 4 and 5, respectively. Support for claims 14 and 15, 16 and 17, and 18-20 may be found in original claims 6-8, respectively. It is believed that these changes do not involve any introduction of new matter, whereby entry is believed to be in order and is respectfully requested.

Respectfully submitted,



Holly D. Kozlowski
Registration No. 30,468
Dinsmore & Shohl LLP
1900 Chemed Center
255 East Fifth Street
Cincinnati, Ohio 45202
(513) 977-8568

VERSION WITH MARKINGS SHOWING CHANGES MADE

In the Claims:

Claims 1-9 are amended as follows:

1. (Amended) Method for increasing the yield of recombinant [proteins] proteins in a microbial fermentation [processes, characterized by the fact that] process, wherein the concentration of [the] a carbon / energy source in the culture is oscillatingly reduced or increased in short cycles.
2. (Amended) Method according to Claim 1, [characterized by the fact that] wherein the oscillations are generated by changing [the] a dosage rate of [the] a feed solution containing the carbon/energy source.
3. (Amended) Method according to [Claims 1 and 2, characterized by the fact that] claim 1, wherein the maximum duration of one cycle is 4 minutes, and the duration of a single phase of the cycle is a maximum of two minutes.
4. (Amended) Method according to [any of Claims 1 through 3, characterized by the fact that] claim 1, wherein the duration of one cycle is one minute, and the duration of a phase of the cycle is a maximum of 75% of the total cycle time.
5. (Amended) Method according to [any of Claims 1 through 3, characterized by the fact that] claim 1, wherein the carbon/energy source is added to the culture in such a manner as to cyclically vary the rate of addition of the substrate solution only during certain segments of the process.

6. (Amended) Method according to [any of Claims 1 through 5, characterized by the fact that] claim 1, wherein the dosage rate is controlled by cyclical activation and deactivation of the addition of the feed solution.

7. (Amended) Method [pursuant to any of Claims 1 through 5, characterized by the fact that] according to claim 1, wherein glucose, glycerol, lactose, galactose, methanol, acetate, molasses, or starch is used as the carbon/energy substrate.

8. (Amended) Method [pursuant to any of Claims 1 through 7, characterized by the fact that,] according to claim 1, wherein, depending on the promoter used, IPTG, [or] indolyl acrylic acid (IAA), [or] lactose, arabinose, galactose, or methanol, [(if not already used as the energy source, is [] are)] added to the culture to induce formation of the recombinant product.

9. (Amended) Method [pursuant to any of Claims 1 through 9, characterized by the fact that] according to claim 1, wherein a temperature shift occurs at the time of the induction of the formation of the recombinant product.